

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**


Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**





Consommation et Corporations Canada	Consumer and Corporate Affairs Canada	(11)	2,009,835
Bureau des brevets	Patent Office	(22)	1990/02/12
Ottawa, Canada		(43)	1990/10/07
K1A 0C9		(52)	260-395
	C.R. CL.		167-245
			167-246
			167-310

5,068,7/99

(51) INTL.CL.<sup>5</sup> A61K-7/48; A61K-31/075; A61K-31/11

(19) (CA) APPLICATION FOR CANADIAN PATENT (12)

(54) Retinal, Derivatives and Their Therapeutic Use

(72) Yu, Ruey J. - U.S.A. ;  
Van Scott, Eugene J. - U.S.A. ;

(73) Same as inventor

(30) (US) 334,518 1989/04/07

(57) 34 Claims

Notice: The specification contained herein as filed

Canada

RETINAL, DERIVATIVES AND THEIR THERAPEUTIC USEAbstract of the Disclosure

Therapeutic as well as preventive measures to improve cosmetic conditions and to alleviate the symptoms of dermatologic disorders with retinal and its derivatives is disclosed. Cosmetic conditions and dermatologic disorders in humans and domestic animals in which retinal and its derivatives may be useful include age spots, wrinkles, warts, eczema, seborrheic keratoses, acne, oily skin, psoriasis, dandruff, xerosis, inflammatory and pruritic skin, disturbed keratinization, skin changes associated with aging and possibly viral infections. Retinal and its derivatives include their stereoisomers, for example, all-trans, 13-cis, 11-cis, 9-cis, 7-cis, 11,13-cis and 9,13-cis vitamin A aldehydes, their hydrate, hemiacetal and acetal forms, and their adduct compounds. Compositions containing retinal or its derivative may be administered systemically such as orally, or topically to the affected areas of the skin.

825-019

RETINAL, DERIVATIVES AND THEIR THERAPEUTIC USE

Be it known that Ruey J. Yu, Ph.D., O.M.D. of 4 Lindenwold Avenue, Ambler, Pennsylvania 19002, and Eugene J. Van Scott, M.D. of 3 Hidden Lane, Abington, Pennsylvania 19001, having made an invention entitled: RETINAL, DERIVATIVES AND THEIR THERAPEUTIC USE.

The following disclosure contains a correct and full description of the invention and of the best mode known to the inventors of taking advantage of same.

Topical treatment with retinal and its derivatives has been found to eradicate or improve age spots, wrinkles, warts, oily skin, psoriasis, eczema, dandruff and other skin changes associated with aging. Oral  
5 administration of retinal has been found to eradicate or improve acne, psoriasis and other disorders of disturbed keratinization and inflammation.

This invention relates generally to therapeutic treatment as well as preventive measures of cosmetic,  
10 dermatologic and other conditions and disorders by either topical or systemic administration of compositions containing retinal or its derivatives. As will be subsequently described in detail, we initially discovered that retinal was effective in the topical  
15 treatment of acne, actinic and nonactinic keratoses.

We have now discovered that compositions containing retinal and its derivatives may have much broader ranges of therapeutic actions on topical or

systemic administration for various cosmetic, dermatologic and other conditions and disorders in humans and animals.

5 In our prior patent application entitled "Process for the Treatment of Acne Vulgaris Utilizing Retinal", Serial No. 458,095, filed April 5, 1974, now U.S. Patent No. 3,932,665, we described and claimed a method of treating acne vulgaris in humans using topical preparations containing retinal. In our patent  
10 application entitled "Treatment of Skin Keratoses with Retinal", Serial No. 723,327, filed September 15, 1976, now U.S. Patent No. 4,034,114, we described and claimed a method of treating actinic or nonactinic keratoses in humans using topical preparations containing retinal.  
15 The actinic keratoses, also known as solar keratoses are found on the sunlight exposed areas of the body such as on the face, head, neck and hands.


#### Description of the Invention

20 It has now been discovered that retinal and its derivatives can be therapeutically useful on topical or systemic administration against varieties of cosmetic and dermatologic conditions and disorders including oily skin, age spots, wrinkles, warts, acne, eczema, seborrheic keratoses, psoriasis, dandruff, xerosis,  
25 inflammatory and pruritic skin, disturbed keratinization, skin changes associated with aging and viral infections.

In accordance with the present invention, retinal and its derivative which are incorporated in  
30 therapeutic compositions for topical, oral or other forms of systemic administration to prevent or alleviate the conditions and symptoms of cosmetic and dermatologic disorders are described as follows.

Retinal, also called vitamin A aldehyde is distinctly different in chemistry from retinol and Retinoic acid. Retinol, also called vitamin A alcohol is commonly known as vitamin A, and is present in fish liver oils as an ester compound, retinyl palmitate. Retinoic acid, also called vitamin A acid is an oxidation product from retinol or retinal. Both retinol and retinal are important in the physiologic functions of vision, growth, reproduction and differentiation, but retinoic acid does not contribute to vision and reproduction in humans and animals.

Retinal can exist in stereoisomeric forms, namely all-trans, 13-cis, 11-cis, 9-cis, 7-cis, 11,13-cis, 9,13-cis. However, the common form is all-trans retinal. Since retinal is chemically an aldehyde it can exist as hemiacetal and acetal forms by reacting with one or two molecules of an alcohol, such as methanol, ethanol or propanol. Such hemiacetal and acetal forms are usually more stable against alkali, and are more resistant to oxidation of the aldehyde group. Retinal may be shown by the following chemical structure:



wherein  $R_1$ ,  $R_2$ =alkyl, aralkyl, aryl or alkoxy group of saturated or unsaturated, straight or branched chain or cyclic form, having 1 to 25 carbon atoms,

$R_3=O$  or  $(OR_4)(OR_5)$ , wherein  $R_4$ ,  $R_5=H$ , alkyl, aralkyl, aryl, diol or polyol group of saturated or unsaturated, straight or branched chain or cyclic form, having 1 to 25 carbon atoms; and the hydrogen atom attached to the carbon atom in the main chain as well as in  $R_1$ ,  $R_2$ ,  $R_4$  and  $R_5$  may be substituted by a halogen atom or a radical such as a lower alkyl, aralkyl, aryl or alkoxy having 1 to 9 carbon atoms. The hemiacetal

and acetal forms are represented by  $-\text{CH}(\text{OR}_4)(\text{OR}_5)$  instead of  $-\text{CHO}$ . The compound may be called retinal hemiacetal when either  $\text{R}_4$  or  $\text{R}_5$  is H, and the compound is called retinal acetal when both  $\text{R}_4$  and  $\text{R}_5$  are for example alkyls or aralkyls. The compound is called retinal hydrate when both  $\text{R}_4$  and  $\text{R}_5$  are H.

Retinal and its derivatives may exist as stereoisomers such as all-trans, 13-cis, 11-cis, 9-cis, 7-cis, 11,13-cis and 9,13-cis forms.

The typical alkyl, aralkyl and aryl groups for  $\text{R}_1$  and  $\text{R}_2$  are for example, methyl, ethyl, propyl, isopropyl, butyl, pentyl, octyl, lauryl, stearyl, benzyl and phenyl etc. The halogen atoms are F, Cl, Br and I. The typical alkoxy groups are methoxy and ethoxy. The typical diol groups are glycol, propylene glycol and 1,3-butanediol. The typical polyol groups include glycerol, butanetriol, inositol and alditols; such as erythritol of tetraol, ribitol of pentaol, mannitol and sorbitol of hexaols.

Retinal or its derivative may react with an unsaturated chemical agent such as maleic anhydride, acetylene dicarboxylic acid or its ester, or hydroquinone to form a crystalline molecular complex called an adduct compound.

The representative retinal and its derivatives which may be useful for topical or systemic administration to improve cosmetic conditions or to alleviate dermatologic disorders are listed below:

1. Retinal;
2. Retinal hydrate;
3. Retinal methyl hemiacetal;
4. Retinal ethyl hemiacetal;
5. Retinal propyl hemiacetal;
6. Retinal isopropyl hemiacetal;



7. Retinal butyl hemiacetal;
8. Retinal pentyl hemiacetal;
9. Retinal octyl hemiacetal;
10. Retinal benzyl hemiacetal;
- 5 11. Retinal dimethyl acetal;
12. Retinal diethyl acetal;
13. Retinal dipropyl acetal;
14. Retinal diisopropyl acetal;
15. Retinal dibutyl acetal;
- 10 16. Retinal dipentyl acetal;
17. Retinal dioctyl acetal;
18. Retinal dibenzyl acetal;
19. Retinal hydroquinone adduct;
20. Retinal maleic anhydride adduct;
- 15 21. Retinal acetylene dicarboxylic acid ester  
adduct;
22. Retinal propylene glycol hemiacetal and  
acetal;
23. Retinal 1,2-O-isopropylidene glyceryl  
20 hemiacetal and acetal;
24. Retinal 3-allyloxy-1, 2-propanediol hemiacetal  
and acetal;
25. Retinal phytyl hemiacetal;
26. Retinal diphytyl acetal;
- 25 27. Retinal dodecyl hemiacetal; and
28. Retinal didodecyl acetal.

Retinal and its derivatives may also be utilized  
in combination with or as additives to enhance  
therapeutic effects of other cosmetic or pharmaceutical  
30 agents to improve cosmetic conditions or alleviate the  
symptoms of dermatologic disorder. Cosmetic and  
pharmaceutical agents include those that improve or  
eradicate age spots, keratoses and wrinkles eradicating  
agents; antiacne agents; antibacterials; antiyeast

agents; antifungal agents; antiviral agents;  
antidandruff agents; antidermatitis agents;  
antipruritic agents; antiinflammatory agents;  
antihyperkeratolytic agents; antidryskin agents;  
5 antipsoriatic agents; antiseborrheic agents; hair  
conditioners and hair treatment agents; antiaging and  
antiwrinkle agents; sunscreen agents; antihistamine  
agents; vitamins; corticosteroids, tanning agents;  
local anesthetics; hormones; retinoids and other  
10 dermatologicals.

Some examples of cosmetic and pharmaceutical  
agents are clotrimazole, miconazole, salicylic acid,  
pramoxine, menthol, retinoic acid, hydrocortisone,  
hydrocortisone valerate, betamethasone valerate,  
15 betamethasone dipropionate, triamcinolone acetonide,  
fluocinonide, hydroquinone, clobetasol propionate,  
benzoyl peroxide, crotamiton, 5-fluorouracil,  
monobenzene, vitamin A palmitate, vitamin E acetate and  
vitamin C.

20 It has been established through tests in humans  
that retinal and its derivatives are therapeutically  
effective for topical, oral, intralesional injection or  
other systemic treatment of various cosmetic conditions  
and dermatologic disorders.

25 Accordingly, it is an object of this invention to  
provide cosmetic as well as medicinal compositions  
containing retinal or its derivatives which when  
topically or systemically administered will  
substantially improve and alleviate the symptoms of  
30 various cosmetic conditions or dermatologic disorders.

It is another object of this invention to provide  
methods for treating various cosmetic conditions or  
dermatologic disorders with topical preparations or  
systemic compositions containing retinal or its

derivative.

#### PREPARATION OF THE THERAPEUTIC COMPOSITIONS

Retinal and its derivatives of the instant invention may be formulated either for topical application or for systemic administration. In the topical preparations retinal and its derivatives may be formulated in aqueous or non-aqueous solution, gel, lotion, cream or ointment containing 0.01 to 5 percent and preferably from 0.01 to 0.5 percent by weight of the total composition. When retinal and its derivatives are formulated in aqueous form, sodium sulfite, sodium bisulfite, sodium metabisulfite or other antioxidants may be added to stabilize retinal and its derivatives in aqueous compositions. Butylated hydroxytoluene (BHT) or butylated hydroxyanisole (BHA), on the other hand, may be added to stabilize retinal and its derivatives in non-aqueous composition. To provide maximal stability of the therapeutic compositions antioxidants of both aqueous and nonaqueous types may also be incorporated into the compositions at the same time. For example, both sodium metabisulfite and BHT may be added to an aqueous acholic solution containing retinal. The concentration of antioxidant may range from 0.01 to 1%.

To prepare a typical aqueous solution, retinal or its derivative is dissolved in a mixture of water, ethanol and propylene glycol in a volume ratio of 30:50:20, respectively. Sodium metabisulfite is then added to the above solution. To prepare a typical non-aqueous solution, retinal or its derivative is dissolved in a mixture of ethanol, isopropyl myristate and squalane in a volume ratio of 70:20:10, respectively. BHT is then added to the above solution.

When a combination composition is desired retinyl palmitate and/or hydroquinone, for example is added to the above non-aqueous solution. The preferred concentration of retinyl palmitate ranges from 1 to 5%.

- 5 The concentration of hydroquinone may range from 1 to 5%, but the preferred concentration is 2% by weight of the total composition.

A typical cream or lotion containing retinal or its derivative is prepared by first dissolving retinal or its derivative in ethanol, acetone, propylene glycol or other solvent. The solution thus prepared is then admixed with commonly available oil-in-water emulsions. BHT or sodium metabisulfite may be added to such emulsions to stabilize retinal or its derivative.

- 15 A typical gel composition is formulated by first dissolving retinal or its derivative in a mixture of ethanol, water and propylene glycol in a volume ratio of 50:30:20, respectively. A gelling agent such as hydroxyethylcellulose, hydroxypropylcellulose or  
20 hydroxypropylmethylcellulose is then added to the mixture with mixing. The preferred concentration of the gelling agent may range from 0.2 to 2 percent by weight of the total composition.

- For oral administration retinal powder is mixed  
25 with gelatin powder so that each gelatin capsule may contain from 5 to 100 mg of retinal. For intralesional injection, retinal or its derivative is dissolved in a vehicle such as isopropyl myristate or safflower oil, and the solution in an injection bottle is sterilized  
30 under standard conditions.

The following are illustrative examples of formulations and compositions according to this invention. Although the examples utilize only selected formulations useful according to this invention, it

should be understood that the following examples are intended to be illustrative and not limiting.

#### EXAMPLE 1

5 All-trans retinal 0.2 g is dissolved in a 100 ml mixture prepared from ethanol, water and propylene glycol in a volume ratio of 50:30:20, respectively. Sodium metabisulfite 0.5 g is added to the mixture with stirring. the light yellowish solution thus formulated contains 0.2% retinal as an active ingredient in  
10 aqueous solution. To prepare a gel composition, hydroxypropyl cellulose 2 g is added to the above solution with mixing until a uniform consistency is obtained.

#### EXAMPLE 2

15 All-trans retinal 0.1 g is dissolved in a 100 ml mixture prepared from ethanol, isopropyl myristate and squalane in a volume ratio of 70:20:10, respectively. BHT 0.1 g is added to the mixture with stirring. The non-aqueous solution thus prepared contains 0.1%  
20 retinal as an active ingredient. To prepare a combination with vitamin A palmitate and hydroquinone, retinyl palmitate 3 g and hydroquinone 2 g are added to the above non-aqueous solution with stirring until a uniform yellowish solution is obtained.

#### EXAMPLE 3

25 A water-washable cream containing retinal 0.1% as an active ingredient may be formulated as follows. 13-cis Retinal 0.1 g is dissolved in 10 ml ethanol, and the solution thus obtained is mixed with 90 g of an  
30 oil-in-water emulsion prepared from petrolatum, glyceryl monostearate, stearyl alcohol, PEG-40

stearate, propylene glycol and water in a weight ratio of 25:5:10:5:10:45, respectively. To prepare a combination with vitamin A palmitate and vitamin E, retinyl palmitate 2 g and alpha tocopheryl acetate 1 g  
5 are added to the above cream with mixing until a uniform light yellowish composition is obtained.

#### EXAMPLE 4

A water-nonwashable lotion containing retinal 0.05% as an active ingredient may be formulated as follows. All-trans retinal 0.05 g is dissolved in 100  
10 g of a lotion prepared from sorbitan trioleate, petrolatum, mineral oil, diisopropyl adipate and isopropyl myristate in a weight ratio of 10:20:10:30:30, respectively. BHA 0.05 g is added to  
15 the mixture with stirring. To prepare a combination with vitamin A palmitate and hydroquinone, retinal palmitate 2 g and hydroquinone 2 g are added to the above lotion with stirring until a uniform light yellowish lotion is obtained.

20

#### EXAMPLE 5

A combination composition containing retinal, hydroquinone and hydrocortisone 17-valerate is formulated as follows. All-trans retinal 0.1 g,  
25 hydroquinone 2 g and hydrocortisone 17-valerate 0.2 g are dissolved in ethanol 10 ml and propylene glycol 5 ml, and the solution thus obtained is mixed with 83 g of commonly available oil-in-water cream containing 0.8% of sodium metabisulfite. Light yellowish cream  
30 thus prepared contains 0.1% retinal, 2% hydroquinone and 0.2% hydrocortisone 17-valerate as active ingredients.

## EXAMPLE 6

A combination composition containing retinal, vitamin A palmitate and benzilic acid may be formulated as follows. All-trans retinal 0.1 g, retinyl palmitate 3 g, benzilic acid 3 g and BHT 0.05 g are dissolved in 94 ml of a mixture prepared from ethanol, isopropyl myristate and squalane in a volume ratio of 60:30:10, respectively. The composition thus formulated contains 0.1% retinal, 3% vitamin A palmitate and 3% benzilic acid. A sun screen agent such as octyl p-methoxycinnamate and a corticosteroid such as betametasone dipropionate may also be incorporated into the above composition.

## EXAMPLE 7

A combination composition containing retinal, gluconolactone, hydroquinone and elastin may be formulated as follows. All-trans retinal 0.05 g, gluconolactone 8 g, hydroquinone 2 g and soluble elastin 0.2 g are dissolved in 90 ml of a mixture prepared from ethanol, water and propylene glycol in a volume ratio of 40:40:20, respectively. Sodium metabisulfite 0.5 g and BHT 0.02 g are added to the above mixture with stirring. The composition thus formulated contains 0.05% retinal, 8% gluconolactone, 2% hydroquinone and 0.2% elastin as active ingredients.

## EXAMPLE 8

A combination composition containing retinal, gluconolactone, hydroquinone, elastin and a sun screen agent may be formulated as follows. All-trans retinal 0.04 g, gluconolactone 5 g, hydroquinone 2 g, soluble elastin 0.1 g and octyl p-methoxycinnamate 3 g are dissolved in 90 ml of a mixture prepared from ethanol,

water and propylene glycol in a volume ratio of 50:30:20, respectively. Sodium metabisulfite 0.4 g and BHA 0.02 g are added to the above mixture with stirring. Concentrated ammonium hydroxide may be added  
5 to adjust the pH to approximately 4.5. The composition thus formulated contains 0.04% retinal, 5% gluconolactone, 2% hydroquinone, 0.1% elastin and 3% octyl p-methoxycinnamate.

## EXAMPLE 9

10 To prepare a composition for intralesional injection retinal 20 mg is dissolved in 10 ml of safflower oil or isopropyl myristate, and the solution thus obtained in an injection bottle is sterilized under standard conditions. The composition thus  
15 prepared contains 0.2% or 2 mg per ml of retinal as an active ingredient.

## EXAMPLE 10

For oral administration gelatin capsules containing retinal or its derivative in different doses  
20 may be prepared as follows. Retinal powder 1 g is thoroughly mixed with 35 g of gelatin powder USP. Each gelatin capsule size No. 0 filled with this mixture contains 10 mg of retinal as an active ingredient. Gelatin capsules containing 25 mg of retinal in each  
25 capsule may also be prepared in the same way but from 1 g of retinal and 14 g of gelatin powder.

## EXAMPLE 11

Retinal dimethyl acetal may be synthesized and formulated in a therapeutic composition as follows.

30 All-trans retinal 100 mg is dissolved in 10 ml of methanol, and concentrated hydrochloric acid 0.04 ml is



added to the mixture with stirring. The reaction mixture changes instantly in color from yellowish to greenish. After 10 minutes at room temperature the mixture is adjusted to alkaline with 0.1 ml of triethanolamine while the color of the solution changes back to yellowish. The mixture is evaporated at 40°C in vacuum to about 2 ml in volume. Ether 10 ml and water 30 ml are added, and the ether layer is separated and washed with 20 ml of water. The ether layer is dried over anhydrous sodium sulfate. After evaporation of ether, yellowish product 95 mg is obtained. Retinal dimethyl acetal thus synthesized is identified by infrared spectroscopy, high performance liquid chromatography and thin-layer chromatograph with a mobility of 0.71 on a solvent system of benzene:methanol 1:1.

Retinal dimethyl acetal 50 mg synthesized from the above procedure is dissolved in 3 ml of acetone, and the solution thus obtained is mixed with 47 g of an oil-in-water emulsion. The composition thus formulated contains 0.1% retinal dimethyl acetal as an active ingredient.

#### EXAMPLE 12

Retinal diethyl acetal may be synthesized and formulated in a therapeutic composition as follows.

All-trans retinal 95 mg is dissolved in 12 ml of ethanol, and concentrated hydrochloric acid 0.04 ml is added to the mixture with stirring. The reaction mixture changes instantly in color from yellowish to brownish. After 30 minutes at room temperature the mixture is adjusted to alkaline with 0.1 ml of triethanolamine while the color of the solution changes back to orange. The mixture is evaporated at 40°C in

vacuum to about 2 ml in volume. Ether 10 ml and water 30 ml are added, and the ether layer is separated and washed with 20 ml of water. The ether layer is dried over anhydrous sodium sulfate. After evaporation of ether, orange product 90 mg is obtained. Retinal diethyl acetal thus synthesized is identified by infrared spectroscopy, high performance liquid chromatography and thin-layer chromatography with a mobility of 0.81 on a solvent system of benzene:methanol 1:1.

Retinal diethyl acetal 50 mg synthesized from the above procedure is dissolved in 3 ml of acetone, and the solution thus obtained is mixed with 30 g of petrolatum and 17 g of mineral oil. The composition thus formulated contains 0.1% retinal diethyl acetal as an active ingredient.

#### EXAMPLE 13

Retinal propylene glycol acetal may be synthesized and formulated in a therapeutic composition as follows.

All-trans retinal 100 mg is dissolved in 10 ml of propylene glycol and 10 ml of ether, and concentrated hydrochloric acid 0.05 ml is added to the mixture with stirring. The reaction mixture changes instantly in color from yellowish to brownish. After 20 minutes at room temperature the mixture is adjusted to alkaline with 0.1 ml of triethanolamine. Ether 10 ml and water 30 ml are added, and the ether layer is separated and washed with 30 ml of water. The ether layer is dried over anhydrous sodium sulfate. After evaporation of ether, orange product 105 mg is obtained. Retinal propylene glycol acetal thus synthesized is identified by infrared spectroscopy, high performance liquid chromatography and thin-layer chromatography with a

mobility of 0.80 on a solvent system of benzene:methanol 1:1.

Retinal propylene glycol acetal 50 mg synthesized from the above procedure is dissolved in 3 ml of acetone, and the solution thus obtained is mixed with 47 g of an oil-in-water emulsion. The composition thus formulated contains 0.1% retinal propylene glycol acetal as an active ingredient.

#### EXAMPLE 14

Retinal 1,2-0-isopropylidene-rac-glyceryl acetal may be synthesized and formulated in a therapeutic composition as follows.

All-trans retinal 100 mg is dissolved in 10 ml of 1,2-0-isopropylidene-rac-glycerol and 5 ml of ether, and concentrated hydrochloric acid 0.02 ml is added to the mixture with stirring. The reaction mixture changes instantly in color from yellowish to orange. After 30 minutes at room temperature the mixture is adjusted to alkaline with 0.1 ml of triethanolamine. Ether 10 ml and water 30 ml are added, and the ether layer is separated and washed with 30 ml of water. The ether layer is dried over anhydrous sodium sulfate. After evaporation of ether, orange product 90 mg is obtained. Retinal 1,2-0-isopropylidene-rac-glyceryl acetal thus synthesized is identified by infrared spectroscopy, high performance liquid chromatography and thin-layer chromatography with a mobility of 0.87 on a solvent system of benzene:methanol 1:1.

Retinal 1,2-0-isopropylidene-rac-glyceryl acetal 50 mg synthesized from the above procedure is dissolved in 3 ml of acetone, and the solution thus obtained is mixed with 47 g of an oil-in-water emulsion. The composition thus formulated contains 0.1% retinal 1,2-

0-isopropylidene-rac-glyceryl acetal as an active ingredient.

#### EXAMPLE 15

5 Retinal 3-allyloxy-1,2-propanediol acetal may be synthesized and formulated in a therapeutic composition as follows.

10 All-trans retinal 100 mg is dissolved in 10 ml of 3-allyloxy-1,2-propanediol and 5 ml of ether, and concentrated hydrochloric acid 0.04 ml is added to the mixture with stirring. The reaction mixture changes instantly in color from yellowish to orange. After 30 minutes at room temperature the mixture is adjusted to alkaline with 0.1 ml of triethanolamine. Ether 10 ml and water 30 ml are added, and the ether layer is separated and washed with 30 ml of water. The ether layer is dried over anhydrous sodium sulfate. After evaporation of ether, orange product 82 mg is obtained. Retinal 3-allyloxy-1,2-propanediol acetal thus synthesized is identified by infrared spectroscopy, 20 high performance liquid chromatography and thin-layer chromatography with a mobility of 0.79 on a solvent system of benzene:methanol 1:1.

25 Retinal 3-allyloxy-1,2-propanediol acetal 50 mg synthesized from the above procedure is dissolved in 3 ml of acetone, and the solution thus obtained is mixed with 30 g of petrolatum and 17 g of mineral oil. The composition thus formulated contains 0.1% retinal 3-allyloxy-1,2-propanediol acetal as an active ingredient.

30

#### EXAMPLE 16

Retinal phytol acetal may be synthesized and formulated in a therapeutic composition as follows.

---

All-trans retinal 284 mg is dissolved in 1 g of phytol and 20 ml of ether, and concentrated hydrochloric acid 0.05 ml is added to the mixture with stirring. The reaction mixture changes instantly in color from yellowish to orange. After 30 minutes at room temperature the mixture is adjusted to alkaline with 0.1 ml of triethanolamine. Water 40 ml is added, and the ether layer is separated and washed with 40 ml of water. The ether layer is dried over anhydrous sodium sulfate. After evaporation of ether, orange product 350 mg is obtained. Retinal phytyl acetal thus synthesized is identified by infrared spectroscopy, high performance liquid chromatography and thin-layer chromatography with a mobility of 0.72 on a solvent system of benzene:methanol 1:1.

Retinal phytyl acetal 50 mg synthesized from the above procedure is dissolved in 3 ml of acetone, and the solution thus obtained is mixed with 47 g of an oil-in-water emulsion. The composition thus formulated contains 0.1% retinal phytyl acetal as an active ingredient.

#### EXAMPLE 17

Retinal dibenzyl acetal may be synthesized and formulated in a therapeutic composition as follows.

All-trans retinal 284 mg is dissolved in 1 g of benzyl alcohol and 20 ml of ether, and concentrated hydrochloric acid 0.04 ml is added to the mixture with stirring. The reaction mixture changes instantly in color from yellowish to orange. After 30 minutes at room temperature the mixture is adjusted to alkaline with 0.1 ml of triethanolamine. Water 40 ml is added, and the ether layer is separated and washed with 40 ml of water. The ether layer is dried over anhydrous sodium

sulfate. After evaporation of ether, orange product 390 mg is obtained. Retinal dibenzyl acetal thus synthesized is identified by infrared spectroscopy, high performance liquid chromatography and thin-layer chromatography with a mobility of 0.84 on a solvent system of benzene:methanol 1:1.

Retinal dibenzyl acetal 50 mg synthesized from the above procedure is dissolved in 3 ml of acetone, and the solution thus obtained is mixed with 47 g of an oil-in-water emulsion. The composition thus formulated contains 0.1% retinal dibenzyl acetal as an active ingredient.

#### EXAMPLE 18

Retinal dipentyl acetal may be synthesized and formulated in a therapeutic composition as follows.

All-trans retinal 284 mg is dissolved in 0.5 g of 1-pentanol and 20 ml of ether, and concentrated hydrochloric acid 0.04 ml is added to the mixture with stirring. The reaction mixture changes instantly in color from yellowish to orange. After 30 minutes at room temperature the mixture is adjusted to alkaline with 0.1 ml of triethanolamine. Water 30 ml is added, and the ether layer is separated and washed with 40 ml of water. The ether layer is dried over anhydrous sodium sulfate. After evaporation of ether, orange product 325 mg is obtained. Retinal dipentyl acetal thus synthesized is identified by infrared spectroscopy, high performance liquid chromatography and thin-layer chromatography with a mobility of 0.83 on a solvent system of benzene:methanol 1:1.

Retinal dipentyl acetal 50 mg synthesized from the above procedure is dissolved in 3 ml of acetone, and the solution thus obtained is mixed with 30 g of

petrolatum and 17 g of mineral oil. The composition thus formulated contains 0.1% retinal dipentyl acetal as an active ingredient.

---

## EXAMPLE 19

Retinal didodecyl acetal may be synthesized and formulated in a therapeutic composition as follows.

5 All-trans retinal 284 mg is dissolved in 500 mg of lauryl alcohol and 20 ml of ether, and concentrated hydrochloric acid 0.05 ml is added to the mixture with stirring. The reaction mixture changes instantly in color from yellowish to orange. After 30 minutes at room temperature the mixture is adjusted to alkaline  
10 with 0.1 ml of triethanolamine. Water 40 ml is added, and the ether layer is separated and washed with 40 ml of water. The ether layer is dried over anhydrous sodium sulfate. After evaporation of ether, orange product 305 mg is obtained. Retinal didodecyl acetal thus  
15 synthesized is identified by infrared spectroscopy, high performance liquid chromatography and thin-layer chromatography with a mobility of 0.78 on a solvent system of benzene:methanol 1:1.

Retinal didodecyl acetal 50 mg synthesized from  
20 the above procedure is dissolved in 3 ml of acetone, and the solution thus obtained is mixed with 47 g of an oil-in-water emulsion. The composition thus formulated contains 0.1% retinal didodecyl acetal as an active ingredient.

## 25 TEST RESULTS

## 1. Animal Studies

Preliminary tests were done on hairless mice and rhino mice. A test composition containing various concentrations of retinal or its derivative was  
30 topically applied to the left dorsal skin of a mouse and a control vehicle was topically applied to the right dorsal skin of the same mouse. Two mice were

---



used for each test composition, and the topical application was carried out once daily, 5 times a week for 2 weeks. At the end of 2 weeks biopsy specimens were taken from the treated sites and also the untreated sites, and were stained and examined histologically. Microscopic examination of the biopsied skin showed that retinal and its derivatives on topical administration increased the number of granular layers, but decreased the thickness of stratum corneum in the skin of the hairless mice and the rhino mice. In the case of rhino mice, retinal and its derivatives had caused comedone impactions of follicular utriculi to disappear, and seemed to diminish the keratinization in the skin of the rhino mice. The skin from the untreated sites and also the vehicle alone treated sites of hairless mice and rhino mice showed no detectable changes under the same conditions.

## 2. Wrinkles and Skin Changes Associated with Aging

Therapeutic compositions containing retinal or its derivative with or without hydroquinone as described in Examples were provided to 23 volunteers and patients having age spots, wrinkles, melasma, lentigines or other pigmented skin spots. Each participating subject received two products, i.e. with or without the addition of hydroquinone to the composition containing retinal or its derivative. The volunteers and patients were instructed to apply topically one medication on one side of the body such as on the back of the left hand and the other medication on the other side of the body such as on the back of the right hand. Specific instructions were given to the participating subjects that the medications were applied twice daily to the

---

skin areas or lesions of age spots. Clinical photos were taken of participating subjects before the initiation of the topical treatment and also at the end of each month after the treatment began.

5        At the end of one month, improvement of age spots was clinically discernible. After two to three months of topical treatment substantial improvement of age spots and fine wrinkles occurred in 19 out of 23 subjects tested. Complete eradication of age spots and  
10       fine wrinkles occurred after two to six months of topical administration with compositions containing retinal or its derivative. Deeper wrinkles seemed to take longer time of topical treatment to show any substantial improvement. Therapeutic compositions  
15       containing both retinal or its derivative and hydroquinone were judged to be more effective in eradicating pigmented age spots, melasma, lentigines and other pigmented skin spots. With the addition of an alpha hydroxyacid such as glycolic acid, lactic acid  
20       and gluconolactone therapeutic compositions containing retinal or its derivative and hydroquinone have been found to be very effective for topical treatment of wrinkles, pigmented or unpigmented age spots, melasma, lentigines and other skin changes associated with  
25       aging.

      Considering various combination compositions containing retinal and other cosmetic and pharmaceutical agents, the therapeutic composition containing retinal, hydroquinone and an alpha  
30       hydroxyacid acid has been found to be one of the most effective formulations for improving wrinkles and the skin changes associated with aging.

### 3. Skin Cleansing and Oily Skin

---

Nine human subjects having oily skin or blemished skin as well as 12 acne patients having extremely oily skin participated in this study. Compositions containing retinal or its derivative may be in a form  
5 of solution, gel, lotion or cream formulation, but most participants preferred a solution or gel over lotion or cream preparation.

Each participating subject received a solution or a gel preparation containing 0.02 to 0.2% of retinal or  
10 its derivative. The participating subjects were instructed to apply topically the solution or gel medication on the affected areas of forehead, face, back and chest. Twice daily administration was continued for 2 to 8 weeks.

15 The degree of improvement of oily skin as well as the rate of improvement of acne lesions were clinically evaluated. Most participants reported that oiliness of skin disappeared within one to two weeks of topical administration, and the skin so treated became smooth  
20 and soft. Two acne patients reported that the solution composition containing 0.2% retinal in water, ethanol and propylene glycol has caused excessive dryness to their skin. However, after they had switched to topical use of 0.05% retinal in a gel preparation the  
25 excessive dryness of skin disappeared, and the acne lesions improved substantially. It was found that 18 out of 21 participants showed substantial improvements on oily skin and acne lesions after eight weeks topical administration of therapeutic compositions containing  
30 retinal or its derivative.

As a skin cleanser for oily skin or acne-prone skin, the therapeutic composition containing retinal or its derivative may also contain alpha hydroxyacids or other cosmetic and pharmaceutical agents. For example,

---

a therapeutic solution or gel composition containing both retinal and benzilic acid has been found to be one of the most effective compositions for oily skin and acne-prone skin.

5      4. Acne: Systemic Administration

Therapeutic compositions containing all-trans retinal for oral administration were formulated as gelatin capsules containing from 10 to 25 mg of retinal as an active ingredient. Four acne patients having  
10      papulo-pustular or cystic acne participated in this study. Each patient was instructed to take orally one capsule each time, three times daily for 2 to 4 weeks.

Two patients having papulo-pustular lesions of acne were taking daily oral dose of 30 to 50 mg of all-trans retinal. The acne lesions improved substantially  
15      after 14 days oral administration of compositions containing all-trans retinal. Two patients having cystic lesions of acne were taking daily oral doses of 75 to 100 mg of all-trans retinal. The acne lesions  
20      improved substantially after 2 to 4 weeks oral administration of compositions containing all-trans retinal.

5. Psoriasis

The involved skin in psoriasis is hyperplastic  
25      (thickened), erythematous (red or inflamed), and has thick adherent scales. The degree of thickening is such that lesions are elevated up to 1 mm above the surface of adjacent normal skin; erythema is usually an intense red; the thickened adherent scales cause the  
30      surface of involved skin to be markedly rough and uneven. These three attributes of thickness, color and texture can be quantified to allow objective

---

measurement of degree of improvement from topically applied test materials or systemically administered medications as follows.

#### Degree Of Improvement

5		None ( 0)	Mild ( 1+)	Moderate ( 2+)	Substantial ( 3+)	Complete ( 4+)
	Thickness	Highly elevated	Detectable reduction	Readily apparent	Barely elevated	Normal thickness
10	Texture	Visibly rough	Palpably rough	Uneven but not rough	Slightly uneven	Visibly and palpably smooth
15	Color	Intense Red	Red	Dark Pink	Light Pink	Normal skin color

By means of such parameters degree of improvement in psoriatic lesions can be numerically recorded and comparisons made of one treated site to another.

#### 20 A. Topical Administration

Therapeutic compositions containing 0.05 to 0.2% retinal or its derivative were formulated as lotion, cream or ointment form. Seven patients having psoriasis participated in this study. Each patient was instructed to apply topically the medication in an amount sufficient to cover circles approximately 4 cm in diameter of involved areas of skin. Applications were made two to three times daily and without occlusive dressings. Generally, the involved skin became less scaly and less erythematous after one week of topical treatment. It was found that five out of

seven patients showed substantial improvement on the treated areas of skin after two weeks topical application of compositions containing retinal or its derivative.

5           B. Systemic Administration

          Compositions containing all-trans retinal for oral administration were formulated as gelatin capsules containing 25 mg retinal as an active ingredient. Two patients having psoriasis participated in this study.  
10   Each patient was instructed to take orally one capsule each time, four times daily for one to two weeks. Both patients shows substantial improvement on psoriatic skin after one week of oral administration. One patient shows a complete clearing of psoriasis lesions  
15   after two weeks oral administration of composition containing all-trans retinal with daily does of 2 mg per kg body weight.

6.   Warts: Intralesional Injections

          Compositions containing 2 mg per ml of all-trans retinal in safflower oil or isopropyl myristate for  
20   intralesional injections were sterilized according to a standard procedure. Wart lesions on the hand and plantar warts on the foot were injected intralesionally with the solution containing all-trans retinal. Each  
25   wart lesion was injected with 0.02 ml solution, i.e. 0.04 mg of all-trans retinal. The same procedure of treatment was repeated on each wart lesion after one week. At the end of 2 weeks from the initial treatment by intralesional injections most wart lesions became  
30   flattened with signs of being eradicated. Some large lesions of warts or more resistant ones showed some signs of improvement after two intralesional

---


injections. It was, however, necessary to repeat the same procedure of treatment a few times more for a complete eradication of wart lesions.

5       The invention may be embodied in other specific  
forms without departing from the spirit or essential  
characteristics thereof. The present embodiments are  
therefore to be considered in all respects as  
illustrative and not restrictive, the scope of the  
invention being indicated by the appended claims and  
10 all changes which come within the meaning and  
equivalency of the claims are therefore intended to be  
embraced therein.

---

Claims

1. A therapeutic composition for cosmetic or dermatologic skin disorders comprising a retinal compound having the formula:



5           wherein

$\text{R}_1$ ,  $\text{R}_2$ =alkyl, aralkyl, aryl or alkoxy group of saturated or unsaturated, isomeric or non-isomeric, straight or branched chain or cyclic form, having 1 to 25 carbon atoms,

10            $\text{R}_3=\text{O}$  or  $(\text{OR}_4)(\text{OR}_5)$ , wherein  $\text{R}_4$ ,  $\text{R}_5=\text{H}$ , alkyl, aralkyl or aryl group of saturated or unsaturated, isomeric or non-isomeric, straight or branched chain or cyclic form, having 1 to 25 carbon atoms; and the hydrogen atom attached to the carbon atom in the main chain as well as in  $\text{R}_1$ ,  $\text{R}_2$ ,  $\text{R}_4$  and  $\text{R}_5$  may be substituted by a halogen atom or a lower alkyl or alkoxy radical having 1 to 9 carbon atoms; or stereoisomers thereof in a pharmaceutically acceptable vehicle for topical application to involved areas or  
15  
20           for internal administration.

2. The composition of Claim 1 wherein said cosmetic and dermatologic disorders include oily skin, dandruff, age spots, warts, eczema, psoriasis, wrinkles, seborrheic keratoses, xerosis, pruritic skin,  
5           disturbed keratinization, inflammatory dermatoses, viral infections and skin changes associated with aging.



3. The retinal compound of Claim 1 wherein said stereoisomers include all-trans, 13-cis, 11-cis, 9-cis, 7-cis, 11,13-cis and 9,13-cis forms.

4. The retinal compound of Claim 1 wherein the compound is a member selected from the group consisting of:

- (1). Retinal;
- 5 (2). Retinal hydrate;
- (3). Retinal methyl hemiacetal;
- (4). Retinal ethyl hemiacetal;
- (5). Retinal propyl hemiacetal;
- (6). Retinal isopropyl hemiacetal;
- 10 (7). Retinal butyl hemiacetal;
- (8). Retinal pentyl hemiacetal;
- (9). Retinal octyl hemiacetal;
- (10). Retinal benzyl hemiacetal;
- (11). Retinal dimethyl acetal;
- 15 (12). Retinal diethyl acetal;
- (13). Retinal dipropyl acetal;
- (14). Retinal diisopropyl acetal;
- (15). Retinal dibutyl acetal;
- (16). Retinal dipentyl acetal;
- 20 (17). Retinal dioctyl acetal;
- (18). Retinal dibenzyl acetal;
- (19). Retinal hydroquinone adduct;
- (20). Retinal maleic anhydride adduct;
- (21). Retinal acetylene dicarboxylic acid ester
- 25 adduct;
- (22). Retinal propylene glycol hemiacetal and acetal;
- (23). Retinal 1,2-O-isopropylidene glyceryl hemiacetal and acetal;
- 30 (24). Retinal 3-allyloxy-1,2-propanediol


hemiacetal and acetal;

- (25). Retinal phytyl hemiacetal;  
(26). Retinal diphytyl acetal;  
(27). Retinal dodecyl hemiacetal; and  
5 (28). Retinal didodecyl acetal.

5. The retinal compound of Claim 1 wherein the compound is a member selected from the group consisting of:

- all-trans Retinal;  
5 13-cis Retinal;  
11-cis Retinal;  
9-cis Retinal;  
7-cis Retinal;  
11,13-cis Retinal; and  
10 9,13-cis Retinal;

6. Method of treatment to improve cosmetic or dermatologic skin disorders administering topically to involved areas of the body or systemically comprising: a retinal compound having the formula:



wherein

$R_1$ ,  $R_2$ =alkyl, aralkyl, aryl or alkoxy group of saturated or unsaturated, isomeric or non-isomeric, straight or branched chain or cyclic form, having 1 to  
10 25 carbon atoms,

$R_3=O$  or  $(OR_4)(OR_5)$ , wherein  $R_4$ ,  $R_5=H$ , alkyl, aralkyl or aryl group of saturated or unsaturated, isomeric or non-isomeric, straight or branched chain or cyclic form, having 1 to 25 carbon atoms; and the  
15 hydrogen atom attached to the carbon atom in the main

chain as well as in  $R_1$ ,  $R_2$ ,  $R_4$  and  $R_5$  may be substituted by a halogen atom or a lower alkyl or alkoxy radical having 1 to 9 carbon atoms; or stereoisomers thereof in a pharmaceutically acceptable vehicle.

7. The method of Claim 6 wherein said cosmetic and dermatologic disorders include oily skin, dandruff, age spots, warts, eczema, psoriasis, wrinkles, seborrheic keratoses, xerosis, pruritic skin, disturbed keratinization, inflammatory dermatoses, viral infections and skin changes associated with aging.

8. The method of Claim 6 wherein said stereoisomers may include all-trans, 13-cis, 11-cis, 9-cis, 7-cis, 11,13-cis and 9,13-cis forms.

9. The method of Claim 6 wherein the compound is a member selected from the group consisting of:


- (1). Retinal;
- (2). Retinal hydrate;
- 5 (3). Retinal methyl hemiacetal;
- (4). Retinal ethyl hemiacetal;
- (5). Retinal propyl hemiacetal;
- (6). Retinal isopropyl hemiacetal;
- (7). Retinal butyl hemiacetal;
- 10 (8). Retinal pentyl hemiacetal;
- (9). Retinal octyl hemiacetal;
- (10). Retinal benzyl hemiacetal;
- (11). Retinal dimethyl acetal;
- (12). Retinal diethyl acetal;
- 15 (13). Retinal dipropyl acetal;
- (14). Retinal diisopropyl acetal;
- (15). Retinal dibutyl acetal;

- 20
- (16). Retinal dipentyl acetal;
  - (17). Retinal dioctyl acetal;
  - (18). Retinal dibenzyl acetal;
  - (19). Retinal hydroquinone adduct;
  - (20). Retinal maleic anhydride adduct; and
  - (21). Retinal acetylene dicarboxylic acid ester adduct.
  - (22). Retinal propylene glycol hemiacetal and acetal;
  - (23). Retinal 1,2-O-isopropylidene glyceryl hemiacetal and acetal;
  - (24). Retinal 3-allyloxy-1, 2-propanediol hemiacetal and acetal;
  - (25). Retinal phytyl hemiacetal;
  - (26). Retinal diphytyl acetal;
  - (27). Retinal dodecyl hemiacetal; and
  - (28). Retinal didodecyl acetal.

10. The method of Claim 6 wherein the compound is a member selected from the group consisting of:

- 5
- all-trans Retinal;
  - 13-cis Retinal;
  - 11-cis Retinal;
  - 9-cis Retinal;
  - 7-cis Retinal;
  - 11,13-cis Retinal;
  - 9,13-cis Retinal;

11. A therapeutic composition for cosmetic or dermatologic skin disorders comprising: in combination a retinal compound having the formula:



5           wherein

$R_1$ ,  $R_2$ =alkyl, aralkyl, aryl or alkoxy group of saturated or unsaturated, isomeric or non-isomeric, straight or branched chain or cyclic form, having 1 to 25 carbon atoms,

10            $R_3=O$  or  $(OR_4)(OR_5)$ , wherein  $R_4$ ,  $R_5=H$ , alkyl, aralkyl or aryl group of saturated or unsaturated, isomeric or non-isomeric, straight or branched chain or cyclic form, having 1 to 25 carbon atoms; and the hydrogen atom attached to the carbon atom in the main  
15 chain as well as in  $R_1$ ,  $R_2$ ,  $R_4$  and  $R_5$  may be substituted by a halogen atom or a lower alkyl or alkoxy radical having 1 to 9 carbon atoms; or stereoisomers thereof; and an effective amount of therapeutic cosmetic or dermatologic agent in a  
20 pharmaceutically acceptable vehicle for systemic administration or for topical application to involved body areas.

          12. The composition of Claim 11 wherein said cosmetic and dermatologic disorders include oily skin, dandruff, age spots, warts, eczema, psoriasis, wrinkles, seborrheic keratoses, xerosis, pruritic skin,  
5 disturbed keratinization, inflammatory dermatoses, viral infections and skin changes associated with aging.

          13. The retinal compound of Claim 11 wherein said stereoisomers include all-trans, 13-cis, 11-cis, 9-cis, 7-cis, 11,13-cis and 9,13-cis form.

          14. The retinal compound of Claim 11 wherein the compound is a member selected from the group consisting of:


- (1). Retinal;
- 5 (2). Retinal hydrate;
- (3). Retinal methyl hemiacetal;
- (4). Retinal ethyl hemiacetal;
- (5). Retinal propyl hemiacetal;
- (6). Retinal isopropyl hemiacetal;
- 10 (7). Retinal butyl hemiacetal;
- (8). Retinal pentyl hemiacetal;
- (9). Retinal octyl hemiacetal;
- (10). Retinal benzyl hemiacetal;
- (11). Retinal dimethyl acetal;
- 15 (12). Retinal diethyl acetal;
- (13). Retinal dipropyl acetal;
- (14). Retinal diisopropyl acetal;
- (15). Retinal dibutyl acetal;
- (16). Retinal dipentyl acetal;
- 20 (17). Retinal dioctyl acetal;
- (18). Retinal dibenzyl acetal;
- (19). Retinal hydroquinone adduct;
- (20). Retinal maleic anhydride adduct;
- (21). Retinal acetylene dicarboxylic acid ester
- 25 adduct;
- (22). Retinal propylene glycol hemiacetal and  
acetal;
- (23). Retinal 1,2-O-isopropylidene glyceryl  
hemiacetal and acetal;
- 30 (24). Retinal 3-allyloxy-1,2-propanediol  
hemiacetal and acetal;
- (25). Retinal phytol hemiacetal;
- (26). Retinal diphytol acetal;
- (27). Retinal dodecyl hemiacetal; and
- 35 (28). Retinal didodecyl acetal.

15. The retinal compound of Claim 11 wherein the compound is a member selected from the group consisting of:

- all-trans Retinal;
- 5 13-cis Retinal;
- 11-cis Retinal;
- 9-cis Retinal;
- 7-cis Retinal;
- 11,13-cis Retinal; and
- 10 9,13-cis Retinal.

16. The composition of Claim 11 wherein said cosmetic and dermatologic agents include antiwrinkle, antikeratosis, antiagespot, antiacne, antidandruff, antipruritic, antibacterial, antifungal, antiyeast, 5 antiviral, antidermatitis, antiinflammatory, antihyperkeratotic, antixerosis, antipsoriasis, antipigmenting and antihistamine agents; and hair conditioners, hair treatment agents, sunscreen agents, local anesthetics, vitamins, corticosteroids, tanning 10 agents, hormones, 5-fluorouracil, retinoids, retinoic acid, retinyl palmitate, monobenzone, hydroquinone, clotrimazole, miconazole, ketoconazole, pramoxine, salicylic acid, menthol, hydrocortisone, hydrocortisone valerate, betamethasone valerate, betamethasone 15 dipropionate, triamcinolone acetonide, fluocinonide, clobetasol propionate, benzoyl peroxide, crotamiton, vitamin E acetate and vitamin C.

17. Method of treatment to improve cosmetic or dermatologic skin disorders comprising administering systemically or topically to involved areas in combination a retinal compound having the formula:



wherein

10             $R_1, R_2$ =alkyl, aralkyl, aryl or alkoxy group of saturated or unsaturated, isomeric or non-isomeric, straight or branched chain or cyclic form, having 1 to 25 carbon atoms,

15             $R_3=O$  or  $(OR_4)(OR_5)$ , wherein  $R_4, R_5=H$ , alkyl, aralkyl or aryl group of saturated or unsaturated, isomeric or non-isomeric, straight or branched chain or cyclic form, having 1 to 25 carbon atoms; and the hydrogen atom attached to the carbon atom in the main chain as well as in  $R_1, R_2, R_4$  and  $R_5$  may be substituted by a halogen atom or a lower alkyl or alkoxy radical having 1 to 9 carbon atoms; or  
20            stereoisomer thereof; and an effective amount of a therapeutic cosmetic or dermatologic agent in a pharmaceutically acceptable vehicle.

18. The method of Claim 17 wherein said cosmetic and dermatologic disorders include oily skin, dandruff, age spots, warts, eczema, psoriasis, wrinkles, seborrheic keratoses, xerosis, pruritic skin, disturbed  
5            keratinization, inflammatory dermatoses, viral infections and skin changes associated with aging.

19. The retinal compound of Claim 17 wherein said stereoisomers include all-trans, 13-cis, 11-cis, 9-cis, 7-cis, 11,13-cis and 9,13-cis form.

20. The retinal compound of Claim 17 wherein the compound is a member selected from the group consisting of:



- (1). Retinal;
- 5 (2). Retinal hydrate;
- (3). Retinal methyl hemiacetal;
- (4). Retinal ethyl hemiacetal;
- (5). Retinal propyl hemiacetal;
- (6). Retinal isopropyl hemiacetal;
- 10 (7). Retinal butyl hemiacetal;
- (8). Retinal pentyl hemiacetal;
- (9). Retinal octyl hemiacetal;
- (10). Retinal benzyl hemiacetal;
- (11). Retinal dimethyl acetal;
- 15 (12). Retinal diethyl acetal;
- (13). Retinal dipropyl acetal;
- (14). Retinal diisopropyl acetal;
- (15). Retinal dibutyl acetal;
- (16). Retinal dipentyl acetal;
- 20 (17). Retinal dioctyl acetal;
- (18). Retinal dibenzyl acetal;
- (19). Retinal hydroquinone adduct;
- (20). Retinal maleic anhydride adduct;
- (21). Retinal acetylene dicarboxylic acid ester
- 25 adduct;
- (22). Retinal propylene glycol hemiacetal and acetal;
- (23). Retinal 1,2-O-isopropylidene glyceryl hemiacetal and acetal;
- 30 (24). Retinal 3-allyloxy-1,2-propanediol hemiacetal and acetal;
- (25). Retinal phytol hemiacetal;
- (26). Retinal diphytol acetal;
- (27). Retinal dodecyl hemiacetal; and
- 35 (28). Retinal didodecyl acetal.

21. The retinal compound of Claim 17 wherein the compound is a member selected from the group consisting of:

- all-trans Retinal;
- 5 13-cis Retinal;
- 11-cis Retinal;
- 9-cis Retinal;
- 7-cis Retinal;
- 11,13-cis Retinal; and
- 10 9,13-cis Retinal.

22. The method of Claim 17 wherein said cosmetic and dermatologic agents include antiwrinkle, antikeratosis, antiagespot, antiacne, antidandruff, antipruritic, antibacterial, antifungal, antiyeast,
- 5 antiviral, antidermatitis, antiinflammatory, antihyperkeratotic, antixerosis, antipsoriasis, antipigmenting and antihistamic agents; and hair conditioners, hair treatment agents, sunscreen agents, local anesthetics, vitamins, corticosteroids, tanning
- 10 agents, hormones, 5-fluorouracil, retinoids, retinoic acid, retinyl palmitate, monobenzone, hydroquinone, clotrimazole, miconazole, ketoconazole, pramoxine, salicylic acid, menthol, hydrocortisone, hydrocortisone valerate, betamethasone valerate, betamethasone
- 15 dipropionate, triamcinolone acetonide, fluocinonide, clobetasol propionate, benzoyl peroxide, crotamiton, vitamin E acetate and vitamin C.

23. The method of Claim 6 wherein said vehicle is a mixture of ethanol, water and propylene glycol with sodium metabisulfite present as a preservative.

24. The vehicle of Claim 23 further comprising hydroxypropyl cellulose present in the mixture to form a gel.

25. The method of Claim 6 wherein said vehicle is a nonaqueous mixture of ethanol, isopropyl myristate and squalane with BHT present as a preservative.

26. The method of Claim 6 wherein said vehicle is a water washable cream comprising a mixture of petrolatum, glyceryl monostearate, stearyl alcohol, PEG-40 stearate, propylene glycol and water.

27. The method of Claim 6 wherein said vehicle is a water nonwashable lotion containing sorbitan trioleate, petrolatum, mineral oil, diisopropyl, adipate and isopropyl myristate with BHA being present as a preservative.

28. The method of Claim 6 wherein said vehicle is gelatin powder in a gelatin capsule for systemic administration.

29. The method of Claim 17 wherein said vehicle is a mixture of ethanol, water and propylene glycol with sodium metabisulfite present as a preservative.

30. The vehicle of Claim 29 further comprising hydroxypropyl cellulose present in the mixture to form a gel.

31. The method of Claim 17 wherein said vehicle is a nonaqueous mixture of ethanol, isopropyl myristate

and squalane with BHT present as a preservative.

32. The method of Claim 17 wherein said vehicle is a water washable cream comprising a mixture of petrolatum, glyceryl monostearate, stearyl alcohol, PEG-40 stearate, propylene glycol and water.

33. The method of Claim 17 wherein said vehicle is a water nonwashable lotion containing sorbitan trioleate, petrolatum, mineral oil, diisopropyl, adipate and isopropyl myristate with BHA being present  
5 as a preservative.

34. The method of Claim 17 wherein said vehicle is gelatin powder in a gelatin capsule for systemic administration.

Smart & Biggar  
Ottawa, Canada  
Patent Agents